

File Revision Date:

October 29, 2024

Data Set Description:

PI: Alain HAUCHECORNE & Fabrice CHANE-MING
Instrument: Rayleigh LIDAR
Site(s): Réunion Island (21.0S, 55.5E, 2158 m)
Measurement Quantities: Temperature (30 - 90 km)

Contact Information:

Name: Alain HAUCHECORNE
LATMOS, UVSQ, 11 Bd d'Alembert
78280 Guyancourt, France
Phone +33 1 80 28 50 24 / +33 6 43 64 94 56
e-mail: alain.hauchecorne@latmos.ipsl.fr

Name: Fabrice CHANE-MING
LACy, Université de la Réunion,
15 avenue René Cassin, Saint-Denis, la Réunion, France
Phone +262 262 93 82 39 / +262 6 92 70 13 27
e-mail fabrice.chane-ming@univ-reunion.fr

Instrument Description:

The same lidar is operated for NDACC temperature and water vapor measurements. Laser pulses are generated by two Quanta Ray Nd:Yag lasers with a repetition rate of 30 Hz, an energy of 375 mJ. per pulse- and a duration of 9 ns. The two lasers are synchronized with a pulse generator with an uncertainty of less than 20 ns. The geometry for transmitter and receiver is coaxial for three reasons: i) to avoid parallax effects, ii) to extend the measurements down to few meters from the ground and iii) to facilitate the alignment. The overlap of the laser beam with the field of view of the telescope is partial from the ground (i.e. 2.2 km asl) to 4 km asl. The backscattered signal is collected by a Newtonian telescope with a primary mirror of 1200 mm diameter (which gives its name to the lidar: "Lidar1200"). No optical fiber is employed: an optical box unit is used directly after the telescope to separate the Raman and Rayleigh signals. The field of view (FOV) of the system is adjustable thanks to a diaphragm field stop at the entrance of the separation unit. A 2 mm FOV (0.3 mrad) allows the background light to be reduced and limits photon counting saturation from low altitude scattering. The separation unit is composed of dichroic beam splitters and interference filters, which separate the backscattered light. Hamamatsu miniature PMT (photomultiplier tubes) and Licel transient recorders are used for the photon detection and data acquisition (in photon counting only). The raw data corresponds to the integration of the signal over 1 minute. In order to reduce saturation effects of the photomultiplier, each channel is equipped with an electronic gating.

Algorithm Description:

The method used to retrieve temperature profiles from molecular backscattered signal and the associated errors has been described in detail by Hauchecorne and Chanin (1980). A description of the

instrumental errors sources and bias has been reported by Keckhut et al. (1993). The two Rayleigh channels are mixed together to provide a single signal for the entire height range. This is achieved in comparing the both channels in the common altitude range (30-50 km) and in calculating the ratio between the both channels and the high-flux non linearity of the high-gain channel considering the low-gain channel as a reference. The signal-induced noise (SIN) is considerably reduced using electronic gating, but still can be identified from the very low mean background noise. It is estimated by fitting with a parabolic function the background signal between 10 km above the top of the temperature profile and 153 km. The residual atmospheric signal at high altitude is estimated using the MSIS model. Computation of temperature profiles requires a pressure initialization. Instead of assuming that the pressure at the top of the profile is equal to the value given by the standard atmosphere model, the scale height of the pressure (which is directly related to the temperature) is adjusting on the MSIS model. Part of the actual algorithm can be found in Keckhut et al. (1993) and in Singh et al. (1996). Data are processed using the V6 version of the Temper code developed by LATMOS. Since the version V4 in 1998 the processing is improved in including in the version V4 an automatic data selection/rejection of data files with too high background signal or too low atmospheric signal (Keckhut et al., 2001).

Expected Precision/Accuracy of Instrument:

The accuracy in determining density and temperature is directly related to photon noise and is associated to temporal and vertical resolution. Statistical noise increases with the altitude and becomes suddenly very large as the signal amplitude reaches the noise level. Relative and absolute uncertainties have been identified and quantified using simulated data (Leblanc et al., 1998). Error calculation can be found in Hauchecorne and Chanin (1980). For NDSC purposes a 2-km vertical resolution constant with altitude is obtained using a Hanning filter. The integration time is about 4 hours, depending on weather conditions. The amplitude of the correction of the non-linearity of the counting is determined with an accuracy of 1 K. The error due to the initialization was estimated to be equal to 15 % at the initialization level. The calculation of uncertainty shows that this error becomes negligible 15 km below as opposed to the noise statistic. The sum of these uncertainties is reported on the NDSC archive. Comparison and data analyses have revealed that the possible bias occurs mainly at the bottom part of the profile induced by mis-alignment problems or by the presence of aerosols.

Instrument History:

A first temperature lidar was operated at Saint-Denis de la Réunion from 1994 to 2009. The new lidar was installed in 2012 in the Maïdo Observatory and the operations started in 2013. In May 2015 a validation campaign took place with the venue of the NASA-GSFC mobile lidar.

Reference Articles:

Limb Temperature Observations in the Stratosphere and Mesosphere Derived From the OMPS Sensor, da Costa P., Keckhut P., Hauchecorne A., Meftah M., Jaross G., Mangin A., Remote Sensing, 2024, 16 (20), pp.3878. (10.3390/rs16203878)

Case Study of a Mesospheric Temperature Inversion over Maïdo Observatory through a Multi-Instrumental Observation, Chane Ming F., Hauchecorne A., Bellisario C., Simoneau P., Keckhut P.,

Trémoulu S., Listowski C., Berthet G., Jégou F., Khaykin S., Tidiga M., et al., *Remote Sensing*, 2023, 15, pp.2045. (10.3390/rs15082045)

Maïdo Observatory: a new high-altitude station facility at Reunion Island (21° S, 55° E) for long-term atmospheric remote sensing and in situ measurements Baray J.-L., Courcoux Y., Keckhut P., Portafaix T., Tulet P., Cammas J.-P., Hauchecorne A., Godin-Beekmann S., De Mazière M., Hermans C. et al, *Atmospheric Measurement Techniques*, 6 (10), 2865-2877, 2013.

Observation of the thermal structure and dynamics of the stratosphere and the mesosphere from space Hauchecorne, A., P. Keckhut, C. Claud, F. Dalaudier, A. Garnier, *Comptes Rendus Geosciences*, 342 (4-5), 323-330, 2010, doi:10.1016/j.crte.2010.01.002

SEASONAL OSCILLATIONS OF MIDDLE ATMOSPHERE TEMPERATURE OBSERVED BY RAYLEIGH LIDARS AND THEIR COMPARISONS WITH TIMED/SABER OBSERVATIONS, X. Dou, T. Li, J. Xu, H. Liu, X. Xue, S. Wang, T. Leblanc, I.S. McDermid, A. Hauchecorne, P. Keckhut, H. Bencherif, G. Heinselman, W. Steinbrecht, M.G. Mlynczak, and J.M. Russell III, *Geophys. Res.*, 2009.

RAYLEIGH LIDAR OBSERVATIONS OF DOUBLE STRATOPAUSE STRUCTURE OVER THREE DIFFERENT NORTHERN HEMISPHERE STATIONS, Sivakumar V., H. Bencherif, A. Hauchecorne, P. Keckhut, D.N. Rao, S. Sharma, H. Chandra, A. Jayaraman and P.B. Rao, *Atmos. Chem. Phys. Discuss.*, 6, 6933-6956, 2007.

AN INSTRUMENTED STATION FOR THE SURVEY OF OZONE AND CLIMATE CHANGE IN THE SOUTHERN TROPICS, JL Baray, J. Leveau, S. Baldy, J. Jouzel, P. Keckhut, G. Bergametti, G. Ancellet, H. Bencherif, B. Cadet, M. Carleer, C. David, M. de Mazière, D. Faduilhe, S. Godin Beekmann, P. Goloub, F. Goutail, J-M. Metzger, B. Morel, J-P. Pommereau, J. Porteneuve, T. Portafaix, F. Posny, L. Robert and M. Van Roozendael, *J. Environ. Monit.*, DOI :10.1039/b607762e, 2006.

STRATOSPHERIC TEMPERATURE MONITORING USING A VIBRATIONAL RAMAN LIDAR. Part 1: AEROSOLS AND OZONE INTERFERENCES, Faduilhe D., P. Keckhut, H. Bencherif, L. Robert and S. Baldy, *J. Environ. Monit.*, 7,357-364, doi:10.1039/B415299a, 2005.

REVIEW OF OZONE AND TEMPERATURE LIDAR VALIDATIONS PERFORMED WITHIN THE FRAMEWORK OF THE NETWORK FOR THE DETECTION OF STRATOSPHERIC CHANGE, P. Keckhut, S. McDermid, D. Swart, T. McGee, S. Godin-Beekmann, A. Adriani, J. Barnes, J-L. Baray, H. Bencherif, H. Claude, G. Fiocco, G. Hansen, A. Hauchecorne, T. Leblanc, C.H. Lee, S. Pal, G. Megie, H. Nakane, R. Neuber, W. Steinbrecht, and J. Thayer, *J. Environ. Monit.*, 6, 721-733, 2004.

EVIDENCE OF TIDAL PERTURBATIONS IN THE MIDDLE ATMOSPHERE OVER SOUTHERN TROPICS AS OBSERVED BY RAYLEIGH LIDAR. Morel B., H. Bencherif, P. Keckhut, S. Baldy, A. Hauchecorne, *J. Atmos. Sol. Terr. Phys.*, 64, 1979-1988, 2002.

ANALYSIS OF GRAVITY WAVE IN THE TROPICAL MIDDLE ATMOSPHERE OVER LA REUNION ISLAND (21°S, 55°E) WITH LIDAR USING WAVELET TECHNIQUES, Chane-Ming F., F. Molinaro, J. Leveau, P. Keckhut et al. Hauchecorne, *Ann. Geophys.*, 18, 485-498, 2000

DESCRIPTION AND EVALUATION OF A TROPOSPHERIC OZONE LIDAR IMPLEMENTED ON AN EXISTING LIDAR IN THE SOUTHERN SUBTROPICS Baray J-L., J. Leveau, J. Porteneuve, G. Ancellet, P. Keckhut, F. Posny et S. Baldy, Appl. Opt., 38, 6808-6817, 1999.

LIDAR DEVELOPMENTS AND OBSERVATIONS OVER REUNION ISLAND (20.8 S, 55.5 E), Bencherif H., J. Leveau, J. Porteneuve, P. Keckhut, A. Hauchecorne, G. MÈgie, F. Fassina and M. Bessafi, Advances in Atmospheric. Remote Sensing, with lidar, Springer-verlag, Eds. : A. Ansmann, R. Neuber, P. Rairoux, and U. Wandinger, 1996.

PLANETARY WAVES AS OBSERVED BY LIDAR OVER REUNION ISLAND, Bencherif H., J. Leveau, A. Hauchecorne, and P. Keckhut, Proceeding of the 1st SPARC General Assembly, Melbourne, December 2-6, 1996.

STRATOSPHERIC TEMPERATURE MEASUREMENTS BY TWO COLLOCATED NDSC LIDARS AT OHP DURING UARS VALIDATION CAMPAIGN, Singh U.N., P. Keckhut, T.J. McGee, Gross, A. Hauchecorne, E.F. Fishbein, J.W. Waters, J.C. Gille, A.E. Roche, and J.M. Russel III, J. Geophys. Res., special issue on UARS Data Validation, 101, 10287-10298, 1996.

MID-LATITUDE LONG-TERM VARIABILITY OF THE MIDDLE ATMOSPHERE TRENDS, AND CYCLIC AND EPISODIC CHANGES, Keckhut P., A. Hauchecorne, and M.L. Chanin, J. Geophys. Res., 100, 18887-18897, 1995.

A CRITICAL REVIEW ON THE DATA BASE ACQUIRED FOR THE LONG TERM SURVEILLANCE OF THE MIDDLE ATMOSPHERE BY FRENCH RAYLEIGH LIDARS, Keckhut P., A. Hauchecorne and M.L. Chanin and A. Hauchecorne, applied Optics, 29, 5182-5186, 1990.

STRATOSPHERE TEMPERATURE MEASUREMENT USING RAMAN LIDAR, Keckhut P., M.L. Chanin and A. Hauchecorne, Applied Optics, 29, 5182-5186, 1990.